



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

August 15, 2016

Richard Kueser  
Regulatory Compliance Director  
Central Solutions, Inc.  
401 Funston Road  
Kansas City, KS 66115

Subject: Label Amendment – Add optional disposal instructions  
Product Name: Low pH Phenolic 256  
EPA Registration Number: 211-62  
Application Date: 05/25/2016  
Decision Number: 518239

Dear Mr. Kueser:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Tom Luminello by phone at (703) 308-8075, or via email at [luminello.tom@epa.gov](mailto:luminello.tom@epa.gov).

Sincerely,



Jacqueline Hardy, Product Manager (34)  
Regulatory Management Branch II  
Antimicrobials Division (7510P)

Enclosure      Stamped Accepted Label dated 8/15/2019

\*\*Medical Device Uses


*(Administrative Note: The following text is required for all labeling, regardless of target market)*

# Low pH Phenolic 256

Germicidal / Detergent Effective in Hard Water  
up to 400 ppm (calculated as CaCO<sub>3</sub>)

With Organic Soil Tolerance

Tuberculocidal - Germicidal - Pseudomonacidal  
Fungicidal - Virucidal\* (Kills HIV-1) - Bactericidal



**ACTIVE INGREDIENTS:**

Ortho-Phenylphenol..... 8.209%

Orthobenzyl-para-Chlorophenol.....6.650%

**INERT INGREDIENTS.....85.141%**

**TOTAL**

**100.00%**

SEE SIDE (REAR) PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**Keep Out of Reach of Children**  
**DANGER**

**Central Solutions, Inc.**

401 Funston Road  
Kansas City, KS 66115

EPA Reg No. 211-62  
EPA Est. No. 211-KS-1

NET CONTENTS: X GALLONS

REV:5/2016

(Side Panel)

*(Administrative Note: The following text is required for all labeling, regardless of target market)***DANGER**

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to by a Poison Control Center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>HOTLINE NUMBER</b> <b>In case of emergency, call your local poison control center or a doctor.</b> <b>Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.</b>	
<b>NOTE TO PHYSICIAN</b> Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.	

Product Questions? Call 1-800-255-0262 during normal business hours (8am-5pm, CST)

**PRECAUTIONARY STATEMENTS****Hazards to Humans and Domestic Animals**

Corrosive. Causes irreversible eye damage or skin burns. Harmful if swallowed or absorbed through skin or inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wear goggles or face shield. Wear protective clothing (long-sleeve shirt and long pants, socks, plus shoed and chemical resistant gloves such as water proof gloves). Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

**USER SAFETY REQUIREMENTS**

Users must wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users must remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Users must remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

*(Administrative Note: The following text is required for labeling destined for poultry, cattle, or swine premise use)***PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Handlers must wear: Coveralls over long-sleeved shirt and long pants; socks and chemical resistant shoes; goggles or face shield and chemical resistant gloves (such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or viton). Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

*(Use Environmental Hazards statement on containers containing ≥ 5 gallons of product):***ENVIRONMENTAL HAZARDS**

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

## **STORAGE AND DISPOSAL**

**Do not contaminate water, food, or feed by storage or disposal.**

**PESTICIDE STORAGE** - Store in a securely closed original container. Store in area that will prevent cross contamination of other pesticides, fertilizer, food, and feed. Avoid storage at temperature extremes or in sunlight.

**PESTICIDE DISPOSAL** - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL** - *(Note to Reviewer: One or more of the following paragraphs for Container Handling will be selected, depending on packaging use/type.)*

Nonrefillable container. Do not reuse or refill this container.

Offer for recycling, if available.

*or*

Offer for reconditioning, if appropriate

*or*

Puncture and dispose of in a sanitary landfill, or by incineration.

### **Less than or equal to 5 gallons:**

Clean container promptly after emptying.

*or*

Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

### **Greater than 5 gallons:**

Clean container promptly after emptying.

*or*

Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

#### ***{For non-refillable SIMPLMIX® 200L Drums}***

Non-Refillable Container. Do not reuse or refill this container. Completely empty container by running pump to dispense all product left in system into application equipment. Dispose of container by permitted onsite incinerator or by transport to offsite permitted incineration services.

If incineration is unavailable, fill the container ¼ full with water. Shake for 10 seconds. Empty and collect rinsate into appropriate storage vessel for disposal. Repeat this procedure two more times. Follow Pesticide Disposal instructions for rinsate disposal. Then offer for recycling or dispose of in a sanitary landfill.

#### ***(For other SimpleMix and F-Style 1 gallons that will be incinerated)***

Completely empty container allowing the product to drain for 10 seconds after the flow begins to drip.

Dispose of container by permitted onsite incinerator or by transport to offsite permitted incineration services.

If incineration is unavailable, fill the container ¼ full with water. Shake for 10 seconds. Empty and collect rinsate into appropriate storage vessel for disposal. Repeat this procedure two more times. Follow Pesticide Disposal instructions for rinsate disposal. Then offer for recycling or dispose of in a sanitary landfill.

***Note to Reviewer: sealed containers are designed to reduce worker exposure to the concentrate. None of these types of containers can be triple rinsed because they are closed, welded, sealed containers.***

*Note to Reviewer: These are representative icons for use sites/application methods that may appear on the label with the appropriate directions for use, PPE or package type.)*



*(Administrative Note: The following text is required for all containers destined for sale and/or distribution in the State of California):*

NOTICE: The State of California has determined that a chemical contained in this product causes cancer based on tests performed on laboratory animals.

*(Administrative Note: The following text is required for labeling destined for the human health care and/or institutional markets.)*

#### **DIRECTION FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### **CLEANING AND DISINFECTION**

For broad spectrum disinfection of gram negative (as demonstrated by efficacy against *Salmonella enterica* and *Pseudomonas aeruginosa*) and gram positive (as demonstrated by efficacy against *Staphylococcus aureus*) bacteria, add ½ ounce LOW pH PHENOLIC 256 per 1 gallon of water. Remove heavy soil or gross filth then apply solution with a mop, cloth, sponge, or hand-pump trigger sprayer so as to wet surface thoroughly. Allow to remain wet for 10 minutes and then air dry. If higher detergency is desired, increase dilution to 1 ounce per gallon of water. This product may be applied only by the methods specified on the labeling.

#### **\*\*Medical Device Uses**

LOW pH PHENOLIC 256 is a general purpose disinfectant that may be used to clean and disinfect hard, non-porous, inanimate, non-critical medical and dental equipment surfaces. LOW pH PHENOLIC 256 may also be used to clean and/or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

*(Administrative Note: The following text is required for labeling on products used to clean and/or disinfect toilet bowls and/or urinals)*

To clean and disinfect toilet bowl and urinal surfaces, remove gross soils from surfaces with bowl brush and flush toilet or urinal prior to applying the use solution. Remove residual bowl water from toilet bowl and urinal by forcing water over the trap with a bowl swab. Press bowl swab against side of the bowl or urinal to remove excess water from the applicator. Add ½ ounce of LOW pH PHENOLIC 256 directly into the bowl or urinal trap water. Swab or brush the interior surface of the fixture thoroughly, making sure to get under the rim. For disinfection, allow to stand for 10 minutes then flush. Wipe exterior surfaces with a cloth or sponge.

*(Administrative Note: The following text is required for labeling destined for the human health care and/or institutional markets. If the customer desires to delete claims against select, specific organisms, it must do so with the approval of the basic registrant.)*

#### **EFFICACY PERFORMANCE**

The broad spectrum efficacy of LOW pH PHENOLIC 256 has been confirmed using A.O.A.C. protocol testing at a use-dilution of 1:256 in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>) in the presence of 5% blood serum (used to simulate organic soil load), at 20°C with an exposure time of 10 minutes.

#### **GERMICIDAL:**

LOW PH PHENOLIC 256 is effective against a broad spectrum of Gram positive and Gram negative pathogens as confirmed by the A.O.A.C. Use-Dilution test, in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>) and in the presence of 5% organic soil load at 20°C with an exposure time of 10 minutes against the following organisms: *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella enterica*, *Methicillin Resistant Staphylococcus aureus (MRSA)*

#### **FUNGICIDAL:**

LOW PH PHENOLIC 256 is effective against the following pathogenic fungi, as confirmed by the A.O.A.C. Use-Dilution test, in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>), and in the presence of 5% organic soil load at 20°C with an exposure time of 10 minutes, in use areas such as shower room floors, locker room benches, bathmats, or any other hard, nonporous, inanimate environmental surfaces: *Trichophyton mentagrophytes* (the athlete's foot fungus)

#### **TUBERCULOCIDAL:**

LOW pH PHENOLIC 256 is effective against *Mycobacterium bovis* (BCG) as demonstrated through testing using the A.O.A.C. Method "Confirmative in Vitro Test for Determining Tuberculocidal Activity", in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>) and in the presence of 5% organic soil load, at 20°C with an exposure time of 10 minutes.

#### **VIRUCIDAL\***

LOW pH PHENOLIC 256 is effective against the following viruses on hard, nonporous, inanimate surfaces, as confirmed by the A.O.A.C. use-dilution test, in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>) and in the presence of 5% organic soil

load, at 20°C with an exposure time of 10 minutes: *HIV-1 (AIDS virus)*, *Vaccina Virus*, *Herpes Simplex virus type 2 (HSV-2)*, *Influenza virus type A<sub>2</sub>*.

Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by influenza A virus. **Low pH Phenolic 256** is a broad-spectrum hard surface disinfectant that has been shown to be effective against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic H1N1 (formerly called swine flu).

or

This product has demonstrated effectiveness against influenza A virus and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 influenza A virus.

or

This product has demonstrated effectiveness against (influenza A virus and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called Swine flu)

or

Kills Pandemic 2009 H1N1 influenza A virus (formerly known as swine flu)

or

Kills Pandemic 2009 H1N1 influenza A virus.

**\*KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type-1 (HIV-1) (associated with AIDS).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** Clean-up must always be done wearing protective latex gloves, gowns, masks, and eye protection.

**CLEANING PROCEDURE:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

**CONTACT TIME:** Leave surface wet for 10 minutes with a ½ ounce per gallon use-dilution (1:256), LOW PH PHENOLIC 256 and water

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood, body fluids, cleaning materials, and clothing must be autoclaved and disposed according to local regulations for infectious waster disposal.

**PELIGRO:** Si no puede leer en ingles, pregunte a su supervisor sobre las instrucciones de su uso apropiadas antes de trabajar con este producto.

*(Administrative Note: The following text is optional for labeling destined for the human health care and/or institutional markets)*

Formulated for Effective Health Care Facility Sanitation

Formulated for Effective Hospital and Health Care Facility Sanitation

Formulated for Health Care Facilities

Formulated for Hospital and Other Health Care Facilities

LOW pH PHENOLIC 256 is a phosphate free, dilutable, hospital germicidal-detergent effective in 10 minutes at 20° C, in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>), in the presence of 5% blood serum, used to simulate organic soil. LOW pH PHENOLIC 256's multi-phenolic formula is designed to clean, disinfect, and deodorize any washable, inanimate, non-porous surface in one easy step. When used as directed, LOW pH PHENOLIC 256 is strong enough to kill a broad spectrum of pathogenic bacteria, yet it is mild enough to have no harmful effects on the surface being disinfected, when used as directed.

LOW pH PHENOLIC 256 has been formulated for use on hard, nonporous, inanimate surfaces commonly found in facilities and areas such as: hospitals, nursing homes, other health care institutions, and establishments that are dedicated to controlling the hazards of cross contamination. This product is specifically designed for use in hospitals, hospital emergency departments, ambulance and patient transfer vehicles, police and law enforcement vehicles, prisoner holding areas, offices, funeral homes, restrooms, medical and dental offices, and day care facilities.

LOW pH PHENOLIC 256 is designed for use on hard, inanimate surfaces such as door handles, medical bed surfaces, springs, wheelchairs, walls, floors, light switches, linen carts, stretcher wheels, hampers, telephones, clean up carts, bassinets, dressing carts, toilet bowl surfaces, urinal surfaces, showers, bathtubs, bed frames, mattress springs, impervious vinyl surfaces, counter tops, table tops, sinks, waste containers, lockers, desks, patient transfer lifts, patient scales, examination tables, dental chairs, medical lamps, stethoscopes, bedpans, blood donor chairs, crutches, instrument trays, operating tables, and other hard, non-porous surfaces that need disinfection.

LOW PH PHENOLIC 256 is a dilutable germicidal detergent formulated for use where broad spectrum disinfection is required in the presence of moderate amounts of organic soil, in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>).

*(Administrative Note: The following text is optional for labeling destined for poultry, cattle, or swine premise use)*

Formulated for Effective Poultry Premise Sanitation  
 Formulated for Effective Swine Premise Sanitation  
 Formulated for Effective Poultry & Swine Premise Sanitation  
 Formulated for Effective Farm Premise Sanitation

*(Administrative Note: The following text is required for labeling destined for poultry, cattle, or swine premise use)*

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

#### **SITE PREPARATION**

The first step in any ongoing sanitation program must be the removal of gross contamination and debris. This may be accomplished using a shovel, broom, or vacuum, depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter. Once the heavy debris is eliminated, disinfection can be accomplished in a single step.

#### **CLEANING AND DISINFECTION**

For all general cleaning and disinfection, thoroughly clean all surfaces with soap and or detergent and rinse with water. Use ½ ounce of LOW pH PHENOLIC 256 per gallon of water (1:256 dilution). Apply LOW pH PHENOLIC 256 using a cloth, mop, or pressure sprayer so as to thoroughly wet surface to be disinfected. Saturate all surfaces with the disinfecting solution; allowing to remain wet for 10 minutes and then let air dry. Immerse halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing litter and manure. This product may be applied only by the methods specified on the labeling.

#### **EFFICACY PERFORMANCE**

The broad spectrum efficacy of LOW pH PHENOLIC 256 has been confirmed using A.O.A.C. protocol testing at a use-dilution of 1:256 in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>), and in the presence of 5% blood serum (used to simulate organic soil load), at 20°C with an exposure time of 10 minutes.

**GERMICIDAL:** LOW pH PHENOLIC 256 is effective against the broad spectrum of Gram positive and Gram negative pathogens including: Staphylococcus aureus, Escherichia coli, Pseudomonas aeruginosa, Salmonella pullorum, Salmonella enterica, Salmonella gallinarum, Mycoplasma synoviae, Bordetella avium, Mycoplasma gallisepticum, Pasteurella multocida, Pseudomonas fragi, Mycobacterium tuberculosis

**FUNGICIDAL:** LOW pH PHENOLIC 256's fungicidal efficacy has been confirmed through testing using the A.O.A.C. Fungicidal Test against the following pathogenic fungi: Trichophyton mentagrophytes, Aspergillus fumigates.

**\*VIRUCIDAL:** Virucidal testing using prescribed EPA testing procedures has verified LOW pH PHENOLIC 256's effectiveness against the following viruses on hard, nonporous, inanimate surfaces, commonly associated with poultry and swine areas including: Avian Influenza Virus, Avian Herpes Virus 2 (Marek's Disease), Avian Bronchitis Virus, Hog Cholera Virus, Avian Laryngotracheitis Virus, Pseudorabies Virus, Avian Adenovirus, African Swine Fever Virus, Avian Reovirus, Infectious Bursal's Disease Virus (IBD)

#### **AREAS OF USE**

**POULTRY PREMISE SANITATION (HATCHERIES):** Egg Receiving Area, Tray Dumping Area, Egg Holding Area, Chick Procession Area, Setter Room, Chick Holding Room or Area, Hatchery Room, Chick Loading Area, Poultry Buildings, Hauling Equipment

#### **APPLICATION AND USE DILUTIONS**

#### **CLEANING AND DISINFECTION**

Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with the disinfectant solution for a period of 10 minutes.

#### **POULTRY HOUSES:**

1. Remove all poultry and feeds from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water.
2. Use ½ ounce of LOW pH PHENOLIC 256 per gallon of water. Saturate surfaces with the disinfecting solution for a period of 10 minutes. Immerse or treat equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing litter and manure. Ventilate buildings, coops, and other closed spaces. Do not house poultry

or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

**HATCHERIES:**

1. Use ½ ounce of LOW pH PHENOLIC 256 per gallon of water to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

**EQUIPMENT:**

1. Empty all troughs, rack, feeding and watering bins, buckets and fountains.  
2. Use ½ ounce of LOW pH PHENOLIC 256 per gallon of water to treat all equipment. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

**VEHICLES:**

1. Clean all vehicles including mats, crates, cabs and wheels with high pressure water and LOW pH PHENOLIC 256 disinfectant. Use ½ ounce per gallon of water to treat all vehicles. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

**SHOEBATHS:**

1. For use in shoebaths: fill shoebath container with LOW pH PHENOLIC 256 diluted at ½ ounce per gallon (1:256). Pre-clean nonporous rubber boots or impervious shoe covers, then expose to the germicidal detergent and allow to remain wet for 10 minutes by air drying.

*(Administrative Note: The following text is required for labeling destined for poultry and/or animal premise use)*

**POULTRY AND ANIMAL INDUSTRY USE**

**FOR CLEANING AND DISINFECTION OF:**

EQUIPMENT including feeders, waterers, fans, racks, troughs, bins and buckets.

VEHICLES including live haul poultry delivery truck, feed trucks, service person cars, tractors, and all other vehicles.

HATCHERIES including floors, walls, equipment, incubators, hatchers, tray and buggies.

SHOEBATH SANITIZERS including all entrances to buildings and hatcheries.

**AREAS OF USE**

**SWINE AND CATTLE SANITATION**

Farrowing Barns and Areas, Waterer & Feeder, Blocks, Nursery, Creep Area, Hauling Equipment

**MISCELLANEOUS**

Horse Stables, Sales Barn Stalls, Dressing Plants, Chutes/Loading Equipment, Sick Pens, Veterinary Hospitals, Cat and Dog Kennels

**GENERAL DISINFECTION USES:**

Can be used in the routine disinfection of hog houses, farrowing houses, finishing houses, growers, feeders and other equipment; horse stables and sick pens at race tracks or sale barns, hospital pens, calf pens and equipment. Acceptable for disinfecting houses used for poultry, turkeys, ducks and game birds and for equipment used in brooding growing and laying quarters. Acceptable for disinfecting dog and cat kennels as well as laboratory animal facilities, at veterinary hospitals, research laboratories and medical institutions. Low pH Phenolic 256 can be used for spraying cattle pens, chutes loading equipment trucks and animal manure pits for the control of odor and ammonia.

**DRESSING PLANT USES:**

Designed for disinfecting equipment, utensils, walls and floors in poultry and animal dressing plants. Equipment and utensils must be thoroughly rinsed with potable water before operations are resumed. Acceptable for disinfecting offal rooms, exterior walls and loading platforms of dressing plants.

**CLEANING AND DISINFECTION**

Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with the disinfectant solution for a period of 10 minutes.

**APPLICATION AND USE DILUTIONS**

**FARM PREMISE:**

DO NOT USE IN MILKING STALLS, MILKING PARLORS OR MILK HOUSES.

1. Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls, and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water.
2. Use ½ ounce of LOW pH PHENOLIC 256 per gallon of water (1 part LOW pH PHENOLIC 256 to 256 parts of water). Saturate all surfaces with the disinfectant solution for a period of 10 minutes.
3. Use 25-50 gallons of solution per 1000 square feet of floor space. Totally immerse equipment to be treated in disinfectant solution. Ventilate building and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or aired. Thoroughly scrub all treated feed racks, troughs, automatic feeder, fountains, and waterers with soap or detergent and rinse with potable water before reuse.

**EQUIPMENT:**

1. Empty all troughs, racks, feeding and watering bins, buckets and fountains.
2. Use ½ ounce of LOW pH PHENOLIC 256 per gallon of water (1 part LOW pH PHENOLIC 256 to 256 parts of water) to saturate all equipment. Allow to stand for 10 minutes then let air dry, or flush and wipe exterior surfaces with a cloth or sponge.

**VEHICLES:**

1. Clean all vehicles including mats, crates, cabs, and wheels with high pressure water and LOW pH PHENOLIC 256 solution.
2. Use ½ ounce of LOW pH PHENOLIC 256 per gallon of water (1 part LOW pH PHENOLIC to 256 parts of water) to treat all vehicles. Allow to stand for 10 minutes then let air dry, or flush and wipe exterior surfaces with a cloth or sponge.

**HATCHERIES:**

1. In the absence of eggs or chicks, clean all surfaces with soap and detergent, and rinse with potable water. Treat all surfaces thoroughly with LOW pH PHENOLIC 256 solution by scrubbing with brush under high water pressure with emphasis on cracks and molding where fungus and mold are usually located.
2. Use ½ ounce of LOW pH PHENOLIC 256 per gallon of water (1 part LOW pH PHENOLIC 256 to 256 parts of water) to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.
3. Do not introduce eggs or chicks until disinfectant solution has been absorbed or dried.
4. Thoroughly scrub treated surfaces with soap or detergent, and rinse with potable water before reuse.

**SHOEBATHS:**

1. For use in shoebaths: fill shoebath container with LOW pH PHENOLIC 256 diluted at ½ ounce per gallon (1:256). Pre-clean nonporous rubber boots or impervious shoe covers, then expose to the germicidal detergent and allow to remain wet for 10 minutes by air drying

*(Administrative Note: The following text is required for labeling destined for poultry and/or animal premise use)*

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**SWINE AND POULTRY BUILDINGS, HATCHERIES, DRESSING PLANTS**

The first step in any sanitation program must be to remove gross contamination and debris. The efficacy of even the most efficient germicide/cleaner is reduced in the presence of heavy organic matter. Remove all animals, poultry, and feeds from buildings, vehicles and enclosures. Empty all troughs, racks and other feeding and watering equipment. Mix LOW pH PHENOLIC 256 at the rate of ½ ounce per gallon of water (1:256) and saturate all surfaces for 10 minutes. Use 25-50 gallons of disinfectant solution per 1000 square feet of floor space. Ventilate buildings and other closed spaces. Do not house livestock poultry or use equipment until treatment has been absorbed set or dried. Rinse equipment with potable water before reuse. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with water before reuse.

In poultry and animal dressing plants, disinfect equipment, utensils, interior and exterior walls, floors, offal and loading platforms with LOW pH PHENOLIC 256 at the 1:256 dilution rate. Again, equipment and utensils must be thoroughly rinsed with potable water before reuse.

**Equipment**

Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Mix ½ ounce of LOW pH PHENOLIC 256 disinfectant per gallon of water (1:256) and treat all equipment for 10 minutes or more. Thoroughly scrub treated feeding and watering appliances with soap or detergent and rinse with potable water before reuse. Clean all vehicles including mats, crates, cabs, and wheels with high pressure water and LOW pH PHENOLIC 256 (1:256 dilution). Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

*(Administrative Note: The following text is optional for labeling destined for the hair salon market)*

LOW pH PHENOLIC 256 is designed for use on brushes, combs, rollers, tweezers, razors, clipper blades, manicure, pedicure, and other salon tools commonly found in beauty salons, barber shops, and hairstyling establishments.

LOW pH PHENOLIC 256 is effective for immersion disinfection of inanimate, non-porous surfaces. Use at a dilution of 1:256 (1/8 ounces of per 32 ounces of water, or 4 ml per liter)

**DISINFECTION:** Thoroughly clean and rinse instruments before immersion in disinfection solution. Immerse instruments completely for 10 minutes at room temperature (20°C/68°F). After immersion, remove instruments and rinse them thoroughly. Fresh solution must be prepared daily or more often if solution becomes visibly dirty.

**CONFIRMATION:** When tested with full immersion at room temperature for 10 minutes in a solution created as above, LOW pH PHENOLIC 256 was confirmed effective against *Mycobacterium tuberculosis*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella enterica*, *Trichophyton mentagrophytes*, *Staphylococcus aureus* (methicillin resistant strain), *Escherichia coli*, and Herpes simplex type 2 on hard, nonporous surfaces

LOW pH PHENOLIC 256 is a dilutable germicidal detergent formulated to clean, disinfect, and deodorize any washable, inanimate, non-porous instrument or surface in one easy step. When used as directed, it is strong enough to kill a broad spectrum of pathogenic bacteria, yet mild enough to have no harmful effects on the surface being disinfected. LOW pH PHENOLIC 256 inhibits mold and mildew, and has a pleasant lavender fragrance.

LOW pH PHENOLIC 256 is acceptable for use on nippers, brushes, combs, rollers, tweezers, razors, clipper blades, manicure, pedicure, and other implements commonly found in salons, including most plastics. When used as directed, this product is formulated to inhibit the development of rust on stainless steel instruments.

*(Administrative Note: The following text is required for labeling destined for pharmaceutical or biotechnology clean room markets. Statements are to be added only to labels of containers destined for those specific markets)*

LOW pH PHENOLIC 256 is acceptable for use by pharmaceutical and biotechnology industries; medical device manufacturers; hospitals and any health care institutions that are dedicated to controlling the hazards of cross contamination in their establishments. For use on hard, inanimate surfaces in aseptic filling and gowning rooms, general manufacturing areas, or on machinery, tables, counters, laminar flow benches, floors, walls, stainless steel, glazed porcelain, glass, and chrome.

**FUNGICIDAL:** At ½ ounce per gallon of water, LOW pH PHENOLIC 256 is also effective against *Trichophyton mentagrophytes* pathogenic fungi in use areas as aseptic filling and gowning rooms, and general manufacturing areas or on any other hard, non-porous, inanimate environmental surfaces.

**VIRUCIDAL\***

LOW pH PHENOLIC 256 is effective against the following viruses as confirmed by the A.O.A.C. use-dilution test, in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>) and in the presence of 5% organic soil load, at 20°C with an exposure time of 10 minutes: *Canine Parvo Virus*, *Feline Parvo Virus*

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

*(Administrative Note: The following text is optional for labeling destined for pharmaceutical or biotechnology clean room markets. Statements are to be added only to labels of containers destined for those specific markets)*

Formulated for Effective Instrument Cleaning and Decontamination Sanitation  
Formulated for Effective Aseptic Controlled Environment ("Clean Room") Sanitation

*(Administrative Note: The following text is optional for the center panel on labeling destined for pharmaceutical or biotechnology clean room markets. Statements are to be added only to labels of containers destined for those specific markets)*

Product and container sterilized and distributed by \_\_\_\_\_.

Sterile pharmaceutical clean room formula.

*(Administrative Note: The following text is optional for labeling destined for the SIMPLEMIX<sup>®</sup> System Technology, closed system dilution mixing system.)*

#### **MARKETING CLAIMS AND STATEMENTS**

SIMPLEMIX<sup>®</sup> System Technology for closed system mixing of pre-measured concentrate and water. *(To be used on applicable container)*

SIMPLEMIX<sup>®</sup> Disinfectant use solution from pre-measured ingredients in this container. Accurate every time. *(To be used on applicable container)*

SIMPLEMIX<sup>®</sup> convenient self-contained mixing system for disinfectant use solution equivalent to 580 ppm active without measuring. *(To be used on applicable container)*

SIMPLEMIX<sup>®</sup> (self-contained bottle/drum design holding pre-measured concentrate/water portions to yield after mixing exact disinfectant use solution of a specified final volume equivalent to ABC ppm active.) *(To be used on applicable container)*

#### **DIRECTIONS FOR USE**

**PREPARATION OF DISINFECTION USE SOLUTION:** For heavily soiled areas in non-medical premise, a preliminary cleaning is required. [(Add) (Mix)] [(1 ounce) (of (this product) (product name))] [(per) (to) (with)] 2 gallons of water (or equivalent use dilution) (solution according to SIMPLEMIX<sup>®</sup> System Container instructions)] (580 ppm active) to disinfect hard non-porous (inanimate) (environmental) surfaces. Apply use solution using a brush, sponge, cloth, mop, auto scrubber, mechanical spray device ((hand pump) coarse pump or trigger spray device) or by immersion so as to thoroughly wet surfaces. For sprayer applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge. Treated surfaces must remain wet for 10 minutes. [(Rinse) (Wipe up excess liquid (with a paper towel))] (Allow to air dry) (Rinsing of floors is not necessary unless they are to be waxed or polished)

(\*\*Rinse all treated surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse). Change cloth, sponge or towels frequently to avoid redeposition of soil. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)].

*(\*\*Note to reviewer: This statement must be used on all labels with food contact surface uses.)*

**PREPARATION OF DISINFECTION/VIRUCIDAL\*/FUNGICIDAL USE SOLUTION:** For heavily soiled areas, a preliminary cleaning is required. [(Add) (Mix)] [(1 ounce) (of (this product) (product name))] [(per) (to) (with)] 2 gallons of water (or equivalent use dilution) (solution according to SIMPLEMIX<sup>®</sup> System Container instructions)] (580 ppm active) to disinfect hard non-porous (inanimate) (environmental) surfaces. Apply use solution using a brush, sponge, cloth, mop, auto scrubber, mechanical spray device ((hand pump) coarse pump or trigger spray device) or by immersion so as to thoroughly wet surfaces. For sprayer applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge. Treated surfaces must remain wet for 10 minutes. [(Rinse) (Wipe up excess liquid (with a paper towel))] (Allow to air dry) (Rinsing of floors is not necessary unless they are to be waxed or polished)

(\*\*Rinse all treated surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse). Change cloth, sponge or towels frequently to avoid redeposition of soil. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)].

*(\*\*Note to reviewer: This statement must be used on all labels with food contact surface uses.)*

**CLEANING AND DISINFECTION:** For heavily soiled areas in non-medical premise, a pre-cleaning step is required. For one-step cleaning and general disinfection, [(Mix) (Prepare) (a)] 1 ounce (of (this product) (product name)) per 2 gallons of water use solution (or equivalent use dilution) (solution according to SIMPLEMIX<sup>®</sup> System Container instructions)] (580 ppm active). Apply use solution to hard nonporous (inanimate) (environmental) surfaces using a brush, sponge, cloth, mop, auto scrubber or mechanical spray device ((hand pump) coarse pump or trigger spray device) or by immersion so as to thoroughly wet surface. For sprayer applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge. Allow surfaces to remain wet for 10 minutes. [(Rinse) (Wipe up excess liquid (with a paper towel))] (Allow to air dry). Change cloth, sponge or towels frequently to avoid redeposition of soil. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)].

**TO DEODORIZE):** For heavily soiled areas, a preliminary cleaning is required. [(Mix) (Prepare) (a)] (1-ounce) (of (this product) (Product Name)) per 2 gallons of water use solution (or equivalent use dilution) (solution according to SIMPLEMIX<sup>®</sup> System Container instructions)] (580 ppm active). Apply use solution to hard, non-porous (inanimate) (environmental) non-food contact surfaces, thoroughly wetting as required. Apply with a brush, cloth, mop, sponge, auto scrubber, mechanical spray device ((hand pump) coarse pump or trigger spray device) or by immersion. For sprayer applications, spray 6-8 inches from surface.

**INSTITUTIONAL DISINFECTANT DIRECTIONS (Hospitals, Dental Offices, Nursing Homes, and Other Health Care Institutions):** For heavily soiled areas, a preliminary cleaning is required. For disinfecting hard, non-porous (inanimate) (environmental) surfaces such as floors, walls, countertops, bathing areas, lavatories, bed frames, tables, chairs, garbage pails, [(add) (mix)] [(1 ounce) (of (this product) (Product Name)) [(to) (with)] 2 gallons of water (580 ppm active) (or equivalent use dilution) (solution according to SIMPLEMIX<sup>®</sup> System Container instructions)] to disinfect hard, non-porous (inanimate) (environmental) surfaces. Apply use solution to surfaces, thoroughly wetting surfaces with a brush, cloth, mop, sponge, auto scrubber, mechanical spray device ((hand pump) coarse pump or trigger spray device) or by immersion. For sprayer applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge. Treated surfaces must remain wet for 10 minutes. [(Rinse) (Wipe up excess liquid (with a paper towel)) (Allow to air dry)]. Change cloth, sponge or towels frequently to avoid redeposition of soil. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)].

**ULTRASONIC BATH DISINFECTANT/FUNGICIDAL DIRECTIONS:** Pre-clean heavily soiled objects. Use ((this product) (Product Name)) to disinfect hard, non-porous (inanimate) (environmental) non-critical objects compatible with Ultrasonic cleaning units. Prepare a fresh solution (of) [(1 ounce) (of (this product) (Product Name)) per 2 gallons of water (or equivalent use dilution) (according to SIMPLEMIX<sup>®</sup> System Container instructions)] (580 ppm active). Pour directly into bath chamber. Place objects into unit and operate for a minimum of 10 minutes (according to manufacturers' use directions). [(Remove objects and [(wash) (rinse)] with sterile water (for injection)) (Allow to air dry)]. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)].

**CLEANING AND DISINFECTING HARD, NON-POROUS (INANIMATE) (ENVIRONMENTAL) SURFACES ON PERSONAL PROTECTIVE EQUIPMENT (RESPIRATORS):** Pre-clean equipment if heavily soiled to ensure proper surface contact. [(Mix) (Prepare) (a)] [(1-ounce) (of (this product) (Product Name)) per 2 gallons of water use solution) (or equivalent use dilution) (solution according to SIMPLEMIX<sup>®</sup> System Container instructions)] (580 ppm active). Apply solution to hard, non-porous (inanimate) (environmental) surfaces of the respirator with a brush, cloth, mop, sponge, auto scrubber, mechanical spray device ((hand pump) coarse pump or trigger spray device) or by immersion. Thoroughly wet all surfaces to be disinfected. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge. Treated surfaces must remain wet for ten (10) minutes. Remove excess solution from equipment prior to storage. Change cloth, sponge or towels frequently to avoid re deposition of soil. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)]. The user must comply with all OSHA regulations for cleaning respiratory protection equipment (29 CFR § 1910.134).

**VETERINARY CLINICS/ANIMAL LIFE SCIENCE LABORATORY/ANIMAL CARE FACILITIES/ANIMAL RESEARCH CENTERS/ANIMAL LABORATORY/ANIMAL QUARANTINE AREAS/ANIMAL BREEDING FACILITIES/ZOOS/PET SHOP/ANIMAL HUSBANDRY/KENNELS/BREEDING AND GROOMING ESTABLISHMENTS/TACK SHOPS DISINFECTION DIRECTIONS:** For cleaning and disinfecting (the following) hard, non-porous (inanimate) (environmental) surfaces ((in the following (premises) (areas to be treated)) ((such as) equipment, utensils, instruments, cages, kennels, stables, stalls and catteries, and other hard-nonporous (inanimate) (environmental) surfaces)), [(saturate with) (use) (apply) (prepare)] [(a use solution of [(1 oz) (of (this product) (Product Name)) per 2 gallons of water) (or equivalent use dilution) (solution according to SIMPLEMIX<sup>®</sup> System Container instructions)] (580 ppm active) (and apply) so as to wet thoroughly. Apply by cloth, mop, brush, sponge, auto scrubber, mechanical spray device ((hand pump) coarse pump or trigger spray device) or by immersion. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge. Allow surfaces to remain wet for a period of 10 minutes. Immerse all halters and other types of hard, non-porous (inanimate) (environmental) equipment used in handling and restraining animals as well as forks, shovels, scrapers

used in removing litter and manure in the use solution. Ventilate buildings and other closed spaces. Do not house (animals) (livestock) or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other treated equipment that dispenses food or water with soap or detergent, and rinse with potable water before reuse. Change cloth, sponge or towels frequently to avoid redeposition of soil. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)].

**SPECIAL INSTRUCTIONS FOR INACTIVATING AVIAN INFLUENZA A (H5N1) VIRUS:** [(Saturate with) (Use) (Apply) (Prepare) (a)] [(the disinfecting solution of) (1 oz) (of (this product) (Product Name)) per 2 gallons of water) (equivalent use dilution) (solution according to SIMPLEMENTIX<sup>®</sup> System Container instructions)] (580 ppm active) (and apply) to all hard nonporous (inanimate) (environmental) surfaces and allow to remain wet for a period of 10 minutes. Apply by cloth, mop, brush, sponge, auto scrubber, mechanical spray device ((hand pump) coarse pump or trigger spray device) or by immersion. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Immerse all halters and other types of hard nonporous (inanimate) (environmental) equipment used in handling and restraining animals as well as forks, shovels, and scrapers used in removing litter and manure. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, and waterers, and other treated equipment with soap or detergent, and rinse with potable water before reuse. Change cloth, sponge or towels frequently to avoid redeposition of soil. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)].

**FUNGICIDAL ACTIVITY:** (Prepare a solution according to SIMPLEMENTIX<sup>®</sup> System Container instructions.) At (this dilution) (1 ounce (of (this product) (Product Name)) per 2 gallons use-level) (or equivalent use dilution) (580 ppm active), [(this product) (Product Name)] is effective against Trichophyton mentagrophytes (athlete's foot fungus) (a cause of Ringworm) (a cause of Ringworm of the foot) on hard nonporous (inanimate) (environmental) surfaces in the presence of 5% blood serum in locker rooms, dressing rooms, shower and bath areas and exercise facilities. Apply use solution by cloth, sponge, mop, brush, auto scrubber, mechanical spray device ((hand pump) coarse pump or trigger spray device). For spray applications, spray 6-8 inches from surface. Do not breathe spray. Saturate surfaces with use solution. Treated surfaces must remain wet for 10 minutes. Allow solution to dry on surface and repeat treatment every seven days or when new growth appears. Change cloth, sponge or towels frequently to avoid redeposition of soil. [(Change the solution) (A fresh solution must be prepared) (Prepare a fresh solution)] at least daily or [(more often if use solution) (when use solution)] becomes visibly [(dirty) (soiled) (diluted)].

**MOLD AND MILDEW CONTROL DIRECTIONS:** (Prepare a solution according to SIMPLEMENTIX<sup>®</sup> System Container instructions.) At (this dilution) (1 ounce (of (this product) (Product Name)) per 2 gallons use-level) (or equivalent use dilution) (580 ppm active), [(this product) (Product Name)] will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, non-porous (inanimate) (environmental) surfaces (as indicated in general instructions above).

**DIRECTIONS FOR ALTERNATE CONTAINERS/DELIVERY SYSTEMS****AUTOMATED DILUTION SYSTEM (DILUTION SYSTEMS TRADE NAME) INSTRUCTIONS:****SIMPLEMIX® SYSTEM CONTAINER:**

*{Trigger Spray Bottle}* lid label:

1. To prepare use solution, open cap.
2. Peel off inner seal by grasping tab at far edge and pulling off.
3. Firmly push small, inner container [(completely) (all the way)] down.
4. Replace cap and tighten.
5. Slowly swirl for 15 seconds.
6. Move spray nozzle to open position.
7. Follow directions for use on label.

or

*{Gallon Size Bottle}* lid label:

1. To prepare use solution, open cap.
2. Peel off inner seal by grasping tab at far edge and pulling off.
3. Firmly push small, inner container [(completely) (all the way)] down.
4. Replace cap and tighten.
5. Slowly swirl for 15 seconds.
6. Open small side spout and peel off inner seal, as above.
7. Pour solution from small side spout onto surfaces to be treated or alternate containers.
8. Follow directions for use on label.

or

*{200 Liter Drum}* lid label

*(Note: To be used with customer supplied pump.):*

1. Close all valves.
2. Uncoil hoses.
3. Connect center hose to pump between X and Y.
4. Open valve 1, then valve 2, then valve 4
5. START pump to empty cubic container
6. When cubic container is empty, turn pump OFF.
7. Close valve 1 and valve 2.
8. Open valve 6 and valve 5.
9. Re-start pump and mix 15 minutes.
10. STOP pump.
11. Close valve 4.
12. To dispense - Open Valves 3 and 7. Run pump only when dispensing.
13. Follow directions for use on label.